

## LIST OF FIGURES

Figure 2.1: Percentage of people with anemia in Andhra Pradesh (NFHS-4) 2015-2016.....	34
Figure 2.2: Percentage of people with anemia in Andhra Pradesh (NFHS-5) 2019-2020 in comparison with NFHS-4 data.....	35
Figure 3.1: Effect of initial metal ion concentration of <i>Lb. fermentum</i> (A) and <i>B. subtilis</i> (B) on biosorption of Fe (II) at pH 4.5, 1 g/L of biomass, 30 <sup>0</sup> C, 24 h, 100 rpm.....	86
Figure 3.2: Effect of biosorbent dose of <i>Lb. fermentum</i> (A) and <i>B. subtilis</i> (B) on biosorption of Fe (II) at 20 mg/L, pH 4.5, 30 <sup>0</sup> C, 24 h, and 100 rpm.....	87
Figure 3.3: Effect of contact time on biosorption of Fe (II) by <i>Lb. fermentum</i> (A) and <i>B. subtilis</i> (B) at 20 mg/L of Fe (II), pH 4.5, 1 g/L biosorbent, 30 <sup>0</sup> C, 100 rpm.....	89
Figure 3.4: SEM image, elemental composition and EDX spectra of unloaded (control) biomass of <i>Lb. fermentum</i> .....	96-97
Figure 3.5: SEM image, elemental composition and EDX spectrum of iron bound <i>Lb. fermentum</i> biomass.....	97
Figure 3.6: SEM image, elemental composition and EDX spectra of unloaded (control) biomass of <i>B. subtilis</i> .....	98
Figure 3.7: SEM image, elemental composition and EDX spectrum of iron bound <i>Lb. fermentum</i> biomass.....	98
Figure 3.8: FT-IR spectra of control (unloaded) <i>Lb. fermentum</i> biomass.....	100
Figure 3.9: FT-IR spectra of Fe (II) bound <i>Lb. fermentum</i> biomass.....	100
Figure 3.10: FT-IR spectra of control (unloaded) <i>B. subtilis</i> biomass.....	101
Figure 3.11: FT-IR spectra of Fe (II) bound <i>B. subtilis</i> biomass.....	102

Figure 3.12: XRD spectra of <i>Lb. fermentum</i> biomass before (A) and after biosorption (B) with Fe (II).....	103
Figure 3.13: XRD spectra of <i>B. subtilis</i> biomass before (A) and after biosorption (B) with Fe (II).....	103
Figure 3.14: Plot of initial pH versus final pH for determination of point zero charge ( $pH_{pzc}$ ) of <i>Lb. fermentum</i> (A) and <i>B. subtilis</i> (B).....	104
Figure 3.15: Phylogenetic relationship of CC30 with different <i>Streptococcus</i> species obtained from BLAST analysis.....	105
Figure 3.16: SE-HPLC chromatogram of CC 30 EPS.....	106
Figure 3.17: GC chromatogram of CC30 EPS.....	106
Figure 4.1: Schematic experimental design of rat feeding experiments.....	112
Figure 4.2: Hemoglobin concentration of standard and experimental treatment groups at different time intervals. Control - CG group fed with SD; ID diet – ID group fed with IDD; ID diet + B + Fe - FT group fed with D1 diet; ID diet (8) + B + Fe (4) – T1 group fed with D1; ID diet (8) + biomass (4) – T2 group fed with D2; ID diet (8) + FeSO <sub>4</sub> .7H <sub>2</sub> O (4) – ST group fed with D3.....	120-121
Figure 4.3: Photomicrographs of the livers of rats fed with SD, IDD, D1, D1, D2 and D3 diets respectively showing no form of necrosis.....	124
Figure 4.4: Photomicrographs of the spleens of rats fed with SD, IDD, FT, D1, D2 and D3 diets, respectively showing no histo-pathological alterations.....	124-125
Figure 5.1: Appearance of chocolate as a result of fortification. R1- control; R2 and R3- free iron fortified chocolate; R4 and R5- iron bound <i>B. subtilis</i> biomass fortified chocolate; R6 and R7- free <i>B. subtilis</i> biomass fortified chocolate.....	133-134
Figure 5.2: Appearance of chocolate as a result of fortification. R1- control; R2 and R3- free iron fortified chocolate; R4 and R5- iron bound <i>Lb. fermentum</i> biomass fortified chocolate; R6 and R7- <i>Lb. fermentum</i> biomass fortified chocolate.....	134-135

Figure 5.3: Appearance of chocolate as a result of fortification. R1- control; R2 and R3- free iron fortified chocolate; R4 and R5- iron uptake *S. cerevisiae* biomass fortified chocolate; R6 and R7- free *S. cerevisiae* biomass fortified chocolate.....135

Figure 5.4: Effect of fortification on TBA (absorbance at 531 nm) of chocolate fortified with *Lb. fermentum* during storage period of 28 days.....137

Figure 5.5: Effect of fortification on TBA (absorbance at 532 nm) of curd/dahi fortified with *Lb. fermentum* during storage period of 7 days.....138

Figure 5.6: Effect of fortification on TBA (absorbance at 531 nm) of chocolate fortified with *B. subtilis* during storage period of 28 days.....138

Figure 5.7: Effect of fortification on TBA (absorbance at 531 nm) of chocolate fortified with *Saccharomyces cerevisiae* during storage period of 30 days.....139